

## The use of *Eucalyptus dives* and Pennyroyal oil in the synthesis of piperitone epoxide (C<sub>10</sub>H<sub>6</sub>O<sub>2</sub>) and P-menthane-8-thiol-3-one (C<sub>10</sub>H<sub>18</sub>OS)

\*Abubakar Sani and Muhammad Sulaiman Rahama

Department of Pure and Industrial Chemistry

Umaru Musa Yar'adua University

Katsina, Nigeria

Corresponding author: Abubakar Sani

---

### ABSTRACT

The aim of this research work is to synthesize piperitone epoxide and p-menthane-8-thiol-3-one, being important constituents of Buchu oil (*Barosma betulina* spp.) from eucalyptus dives oil and pennyroyal oil. 10 mL of eucalyptus dives oil was treated and shaken well with excessive saturated sodium hydrogen sulfite for an hour, which produced a white precipitate of complex mixture of aldehydes and ketones. The mixture further undergoes several treatments with chemicals and finally piperitone was isolated via fractional distillation at temperature range between 94-96 °C. Furthermore, Piperitone Epoxide and P-menthane-8-thiol-3-one were synthesized from piperitone and pennyroyal oil respectively. The synthesized compounds were characterized using Thermogravimetric analysis (TGA) and Attenuated total reflection Fourier trans-infrared (ATR-FTIR) analyses. The results obtained showed the inflection point of pennyroyal oil (157 °C) falls within the range of inflection point of most the essential oils such Eugenol (162 °C), cinnamon (156 °C) etc. In the infrared analysis slight shift of structural bonds was observed, which indicates chemical reactions and synthesis re-arrangement of reacting species occurred. In conclusion, the compounds proposed to be synthesized were successfully synthesized and characterized.

**KEYWORDS:** *Eucalyptus dives* oil, pennyroyal oil, piperitone epoxide, p-menthane-8-thiol-3-one, Infrared spectroscopy.

---

Date of Submission: 20-08-2020

Date of acceptance: 06-09-2020

---

### I. INTRODUCTION

The term essential oils derived its name as an extract from the characteristics fragrance of the plant from which is obtained. They are complex aromatic, volatile compositions oils of obtained from a plant source or material either by steam or dry distillation, using a suitable heating or mechanical method of extraction and usually carry the odour of their plant [1]. In addition, essential oils being aromatic in nature are basically extracted from aromatic plant sources, mostly biosynthesis by the plant organs as secondary intermediate products. The aromatic plants make only 10% of over 1700 plant species globally and the genres can be traced in a small number of families, which includes Asteraceae, Rutaceae, Piperaceae Lauraceae, Lamiaceae, Myrtaceae, cupressaceae, and Poaceae. The essential oils as a volatile complex mixture of compounds obtained from aromatic plant extracts can be categorised into two distinct groups based on the chemical composition. These includes hydrocarbons compounds which are made up of terpenes (isoprenes), terpenoids that is subdivided into monoterpenes, sesquiterpenes, diterpenes and the oxygenated compounds which consist of mainly aldehydes, ketones, alcohols, phenol and oxides [1, 2]. The isoprene unit is also the basic building block of terpenes with general formula of (C<sub>5</sub>H<sub>8</sub>)<sub>n</sub>, where 'n' represents the number of attached isoprene units. The linkage system of isoprene units is in head-to-tail arrangement [2].

The essential oils been products of secondary metabolites of aromatic plants are composed of complex mixture of volatile constituents that are used in flavourings, perfumes, insects and animal repellents. They are also used in cosmetics and aromatherapy applications as an anti-inflammatory, anti-septic, anti-viral and anti-bacterial [3]. The essential oils can be served as food supplements for lactating animals as well as food preservatives [4]. These oils are mainly extracted from different plant sources.

Buchu oil, an essential oil mainly produced from Buchu plant, consist of many compounds. Buchu plant is extensively used as tonic and medicinal stimulant. Among the main chemical constituents found in the *Agathosma betulina* specie are limonene, cis mercaptone, trans mercaptone, 1-pulegone, piperitone, piperitone epoxide, diosphenol, d-pulegone, 4-diosphenol, iso-menthone and menthone. In the specie of *Barosma*

*crenulata*, significant amount of l-pulegone, P-menthane-8-thiol-3-one, limonene, cis acetylthiol, trans acetylthiol, cis mercaptone, trans mercaptone, menthone and trace of diosphenol are found. The l-pulegone is exerting toxicity effect, if consumed in large quantities [5]. The essential oil obtained from *Eucalyptus dives* tree is usually produced by steam distillation, having two chemotype of piperitone and cineole. The piperitone chemotype of *Eucalyptus dives* (53% composition of piperitone) has a peppermint aroma and flavor and can be used in the production and synthesis of menthol, piperitone, thymol etc. while the cineole chemotype of *Eucalyptus dives* is uncommon in nature and rich in with cineole about 70-80% composition [6]. The dominant chemical compositions in *Eucalyptus dives* (*E. dives*) essential oil are (40.5%) piperitone, (17.4%) phellandrene, (8.5%) para-cymene and (4.7%) terpin-4-ol [7]. The distillate heavy product of the *Eucalyptus dives* specie found in South Africa contains significant amount piperitone, while the distillate light product contains high quantity of  $\alpha$ -thujen,  $\alpha$ -terpinolene and  $\alpha$ -phellandrene with several other volatiles. The Australian specie of *Eucalyptus dive* has a unique advantage of high piperitone content extracts (42.9%) and  $\alpha$ -phellandrene (30.0%) as the dominant constituents in the essential oil. Current attention and interest have been now focusing on the synthetic chemistry of their functional group properties to synthesize numerous compounds of commercial value such as menthol, piperitone epoxide etc. The *E. dives* essential oil has different colour ranges from colourless when refined to pale yellow when first obtained by steam fractional distillation of the leaves. The *E.dives* essential oil possesses wide applications in chemical purposes and pharmaceutical formulation. This includes insect repellent, antiseptic, flavour in in food and fragrance manufacture, and as a starting material or component for industrial compounds. It also showed antimicrobial activity against yeast and other bacterial infection, which can be used in cosmetic formulation.

Pennyroyal popularly known as *Mentha pulegium* is an aromatic flowering plant that belongs to the family lamiaceae. Its volatile essential oil obtained by steam distillation or hydro-distillation of *Mentha pulegium* L. (European pennyroyal specie) and *Hedoema pulegiodes* L. (American pennyroyal specie), basically from the leaves and flowering tops. The crushed leaves of pennyroyal produced a strong aroma of almost similar to spearmint smell and the pungent odour of it essential oil is related to the high composition of pulegone which is toxic organic compound [8]. According to the Petrakis et al [80], the main constituents of pennyroyal essential oil consists of pulegone (61.3%- 77.9%), iso-menthone (10.6%-18.5%), piperitone (0.3%-3.2%), menthone (0.6%-8.3%) and cis-iso pulegone (0-1.7%), while another analysis states that, the *Mentha pulegium* (European origin) essential oil is made up of 53 different constituents, making up of 86.3% of the oil composition.

This work would focus on the synthesis and characterization of piperitone epoxide and p-menthane-8-thiol-3-one, being important constituents of Buchu oil (*Barosma betulina spp.*) from eucalyptus dives oil and pennyroyal oil. These compounds have economic values in terms of wider application in traditional and modern medicinal science and flavours, fragrances and cosmetics usage. The commercial high demand of these compounds exposes the plant to the extinction threat, which motivates scientist to initiate synthetic routes to manufacture major constituents in the Buchu oil.

## II. MATERIALS AND METHODS

### Reagents

The solvents and reagents used in the research work includes sodium hydroxide (Reagent grade,  $\geq$  98%, pellets (anhydrous) sigma-aldrich), sodium hydrogen sulfite (Purum, 40% sigma-aldrich), diethylamine ( $\geq$  99.5% sigma-aldrich), sodium sulfate (ACS reagent,  $\geq$  99% anhydrous, granular, sigma-aldrich), 30% hydrogen peroxide (30wt,% in H<sub>2</sub>O reagent, sigma-aldrich), methanol (CHROMA SOLV® Plus, for HPLC,  $\geq$  99.9% sigma-aldrich), acetone (CHROMA SOLV® Plus, for HPLC,  $\geq$  99.9% sigma-aldrich), n-hexane (CHROMA SOLV® Plus, for HPLC,  $\geq$  97.0% (GC) sigma-aldrich), hydrochloric acid (ACS reagent, 37% sigma-aldrich), diethyl ether (ACS reagent, anhydrous,  $\geq$  99.7% sigma-aldrich), iron (ii) sulfide (technical grade, sigma-aldrich), Pennyroyal oil ( $\geq$  90% aldrich) and *Eucalyptus dives* oil ( $\geq$  90% aldrich). All reagents are purchased and supplied by Sigma Aldrich chemical company Inc.

### Apparatus

Three neck 100 ml round bottom flask, two neck 100 ml round bottom flask, fractionating column, condenser, thermometers, beakers, separating funnel, dropping(addition) funnel, adaptors, funnel, magnetic stirrer, vials, rotatory evaporator etc.

### Piperitone Purification from *Eucalyptus dives* oil

The purification procedure consisted of reaction stages, whereby 10ml of *Eucalyptus dives* oil was treated and well shaken with excessive saturated sodium hydrogen sulfite for an hour, which produced a white precipitate of complex mixture of aldehydes and ketones. The mixture was treated and washed with portions of methanol and diethyl ether for the purification. The solution obtained was dissolved in hot water and a few

drops of 1M of sodium hydroxide were added to regenerate ketone compounds before drying with sodium sulfate. Fractional distillation was setup to isolate piperitone at temperature range between 94-96 °C.

### **Piperitone Epoxide**

The experiment involved an epoxidation reaction of pure piperitone obtained from experiment 1. 5ml of pure piperitone was treated with 10ml of 30% Hydrogen Peroxide (H<sub>2</sub>O<sub>2</sub>) in 16ml methanol at -15 °C. Furthermore, 3ml of 6M aqueous sodium hydroxide was added with stirring ensuring the temperature of the reaction did not exceed 0 °C. After the addition was complete, the mixture was continued stirring for further two hours. 20ml of cold water was added to the mixture and extracted four times with portion of diethyl ether (4×12ml). The extract was also washed with water and dried using anhydrous sodium sulfate. A small quantity of diethyl ether remaining in the extract was removed using a rotary evaporator at 32 °C and the Piperitone epoxide was distilled at 105 °C by fractional distillation.

### **P-menthane-8-thiol-3-one**

This is an experiment involved the synthesis of organosulfur compound (p-menthane-8-thiol-3-one) from d-pulegone obtained from fractional distillation of pennyroyal oil. The synthesis processes includes the immersion of 5ml of d-pulegone into 5ml diethylamine in a cooling bath. A continued flow of hydrogen sulfite gas was bubbled into the reaction mixture for 5 hours and the temperature of reaction did not exceed 30 °C. A crystalline residue was obtained, which was acidified with hydrochloric acid and extracted two times with portion of ether (2×10ml). The extract was washed with water and dried with anhydrous sodium sulfate. A small quantity of diethyl ether remaining was removed using rotatory evaporator from the extract and a yellow oil of thiol compound is obtained.

### **Instrumental Analysis**

Thermogravimetric analysis (TGA) was carried out on the starting materials (i.e. essential oils) to provide a quantitative measurement of weight loss changes associated with thermally induced transitions. The technique was necessary due to fractional distillation process in the research work that is conducted in isothermal conditions. The TGA operating parameters are as follows:

Mass of sample: 7.0593 mg of pennyroyal oil

Start temperature: Ambient

Heating rate: 10 °C min<sup>-1</sup>

Atmosphere: Nitrogen 20ml min<sup>-1</sup>

End temperature: 500 °C


Pan: Platinum

For each analysis, 7.0593 mg of the sample was measured in the sample pan and transferred into the instrument for the analysis under the above set of parameters. A thermogram was obtained after the analysis, showing a derivative curve of mass change against temperature change.

Gas chromatography-Mass spectrometry (GC-MS) was used to identify the components present in the oil using an Agilent Technologies 6890N GC System with an ID-BPX5 column (30 m × 0.25 mm × 0.25 μm). The electron energy and source temperatures were 70 eV and 280 °C respectively. The GC was operated under an injection split mode.


Attenuated total reflection Fourier trans-infrared (ATR-FTR) spectroscopy was used to analyse the surface attached molecules and determine their secondary structural pattern, orientation and interaction. The analysis of these compounds was conducted under following conditions: scanning times of 8, resolution 4 cm<sup>-1</sup> and scanning frequencies ranges from 400-4000 cm<sup>-1</sup>. The background spectrum was carried out before each of the samples was placed on the attenuated total reflectance cell provided with a diamond crystal. The spectral curves were recorded using OMNIC software programmed in the instrument.

### III. RESULTS AND DISCUSSIONS




**Fig.1. Infrared spectrum of Eucalyptus dives oil**

Fig.1 above shows the infrared spectrum of Eucalyptus dives oil. It revealed two strongly absorbing peaks at about 3000 and 1700  $\text{cm}^{-1}$  for C-H and C=O stretching frequencies respectively. The intense peak at around 2957.86  $\text{cm}^{-1}$  corresponds to C-H stretch (aromatic) and the peak at around 2870.69  $\text{cm}^{-1}$  is assigned to methyl and methylene stretching. The strong absorption at 1668.02  $\text{cm}^{-1}$  corresponds to carbonyl group (C=O) in piperitone, which is intense due to the strong absorption of infrared radiation by the C=O bond. The absorbance at 1637.54  $\text{cm}^{-1}$  is assigned to alkene (C=C bond) and also the peak at around 1431.46 and 1366.88  $\text{cm}^{-1}$  corresponds to C-H bend and CH<sub>3</sub> bend respectively. Aromatic stretch (out-of-plane bend) is observed at 815.66 – 890.71  $\text{cm}^{-1}$ .




**Fig.2. Infrared spectrum of pure piperitone**

The pure piperitone spectrum exhibited similar characteristic to the Eucalyptus dives oil IR spectrum in terms of structural information and frequencies at which various functional group absorbed infrared radiation as highlighted in Fig. 2. The significant differences is observed in the intense absorption bands of C-H stretch (aromatic) at 2958.45  $\text{cm}^{-1}$  and slightly shift of carbonyl group (C=O) from 1668.02  $\text{cm}^{-1}$  in Eucalyptus dives oil to 1671.78  $\text{cm}^{-1}$  for the pure piperitone, which is within the carbonyl group frequencies range (1850 – 1630  $\text{cm}^{-1}$ ).




**Fig.3. Infrared spectrum of Piperitone epoxide**

Fig. 3 showed the spectrum of piperitone epoxide. The spectrum showed two intense absorbing peaks and numerous peaks in the fingerprint region. The strong absorption peak at  $2959.37\text{ cm}^{-1}$  is assigned to C-H stretch (aromatic) and the band at  $2930.26\text{ cm}^{-1}$  corresponds to methyl stretching. The strong peak observed at  $1701.75\text{ cm}^{-1}$  is assigned to unconjugated carbonyl group (C=O). The distinct weak and strong peaks at  $1278.10$  and  $798.40\text{ cm}^{-1}$  in the fingerprint region were assigned to ring-stretch and asymmetric ring deformation of the epoxide ring respectively.




**Fig.4. Infrared spectrum of pulegone**

As shown in Fig.4 above, the pulegone spectrum showed two strongly absorbing peaks at about  $3000$  and  $1700\text{ cm}^{-1}$  for C-H (aromatic) and C=O stretching frequencies respectively. The intense peak at around  $2953.85\text{ cm}^{-1}$  corresponds to C-H stretch (aromatic) and the peak at around  $2870.69\text{ cm}^{-1}$  is assigned to methyl and methylene stretching. The strong absorption at  $1668.02\text{ cm}^{-1}$  corresponds to the carbonyl group (C=O) in pulegone, which is intense due to strong absorption of infrared radiation by C=O bond. The absorbance at  $1618.64\text{ cm}^{-1}$  is assigned to alkene (C=C bond) and also the peak at around  $1455.07$  and  $1363.45\text{ cm}^{-1}$  corresponds to C-H bend and  $\text{CH}_3$  bend respectively. Aromatic stretch (out-of-plane bend) is observed at  $645.51 - 875.54\text{ cm}^{-1}$ .



**Fig.5. Infrared spectrum of P-menthane-8-thiol-3-one**

The spectrum shown in Fig.5 above revealed dominated absorption peaks characteristics of C-H stretch at 2953.59 cm<sup>-1</sup> and C=O stretch at 1708.60 cm<sup>-1</sup>. The weak absorption band of S-H stretch at 2361.90 cm<sup>-1</sup> is observed, which corresponds to the thiol group. The absorbance at 1618.57 cm<sup>-1</sup> is assigned to the alkene (C=C bond) and also peak at around 1455.20 and 1366.65 cm<sup>-1</sup> corresponds to C-H bend and CH<sub>3</sub> bend respectively. Aromatic stretch (out-of-plane bend) is observed at 645.57 – 875.87 cm<sup>-1</sup>.



**Fig.6. TGA curve of Pennyroyal oil sample**

Fig.6 above reveals the TGA analysis of pennyroyal oil. The pennyroyal oil was heated from 35.55 °C to 500 °C (at heating rate 10 °C min<sup>-1</sup>) in a dry nitrogen atmosphere, flowing at 20ml min<sup>-1</sup>. The curve obtained in nitrogen atmosphere has shown starting decomposition at temperature of 140.40 °C and the weight loss process ended at 215.18 °C. The TGA curve showed the endothermic curve of the analysis, which reflects the characteristic by pointing from maximum value to the baseline as weight loss with temperature increases, which exhibits zero order kinetics. The inflection point of pennyroyal oil obtained from the TGA curve of 157.05 °C is slightly low compare to other essential oils such as long pepper of inflection point of 182.05 °C [31], similar to cinnamaldehyde (163 °C), Eugenol (162 °C) higher than cineole (101 °C) and almost the same with cinnamon oil (156 °C) [10].

### Reaction Mechanism for Piperitone purification

The piperitone purification involved the nucleophilic addition of bisulfite to form a chemical bond [11]. The plane polarized nature of C=O group makes the carbon atom partial charge and attracts negatively charged nucleophile.

In the double bond of carbon-oxygen atoms, the electrons are not shared equally. The carbon atom has partial positive charge and the pi electron of the carbonyl bond moves toward to the oxygen atom which is more electronegative than the carbon; therefore, the carbon atom has a strong tendency to attract electrons to itself in a carbon-oxygen bond. This causes the carbon-oxygen double bond to be highly polar in a nature [12].

The bisulfite ion from the sodium hydrogen sulfite produced a weak acid and covalent molecule that dissociates and loses a hydrogen ion, leaving an electron behind on the sulfite ion, with negative charge.

The carbonyl double bond is made up of  $\sigma$  and  $\pi$  bond. The  $\sigma$  bonds between carbon and oxygen hybridized atoms produced two  $sp^2$  orbitals. The  $sp^2$  orbitals on the carbon atom form two  $\sigma$  bonds to the substituents and the other  $sp^2$  orbitals of the oxygen are filled by two lone pairs. This makes the carbonyl group to be plane polarized in nature [12].

The Nucleophilic addition reaction occurred between the sulfite ions with the positive carbon atom to formed single bond, while the negative ion of the oxygen atom removed a hydrogen ion from the bisulfite molecule to form another single bond respectively. The sodium hydroxide added in the reaction regenerate piperitone in its pure form by reversing the mechanism. About 7 ml of pure piperitone was obtained after the fractional distillation.

#### **Reaction Mechanism for Piperitone epoxide**

This reaction involves methanol deprotonation of hydrogen peroxide (good oxidizing agent), which provides an active site for  $Na^+$  attached to the peroxide forming a conjugate base of hydrogen peroxide (acid-base reaction). The  $Na^+$  arises from the dissociation of sodium hydroxide in the reaction. Methanol acts as a nucleophile in the reaction, thereby establishing a nucleophilic attack on the electron-deficient double bonds of piperitone by donating a pair of protons, forming a C-H bond (exothermic reaction). This protonation converts the carbon atom into carbonium ion (reactive carbon) which becomes susceptible to nucleophile attack of conjugated base of peroxide. A recycled methanol molecule is obtained after this reaction.

Asymmetric carbon atom is formed from the nucleophilic attack on the double bonds, which features the causes of chirality in the new compound to be formed (Piperitone epoxide). An asymmetric carbon atom is carbon atom attached to four different group atoms or atoms [13].

A weak single bond in the conjugated base of hydrogen peroxide (a better nucleophile) dissociate into  $Na^+$  and  $^{\ominus}OOH$  radicals, attacked the carbocation of piperitone compound either from above or below of the bond. This sets up migration reactions, in which the asymmetric carbon atom released a hydrogen ion and reacts with hydroxide ion of peroxide, while hydroxide ion and sodium ion formed sodium hydroxide.

Two optical isomers of "R" and "S" piperitone epoxide are expected at the end of the reaction (enantiomers) due to asymmetrical double bonds reaction.

#### **Reaction Mechanism for P-menthane-8-thiol-3-one**

Diethylamine is a good nucleophile for conjugation addition reactions, possessing nitrogen atom with an unshared electron pair that act as nucleophiles toward the alkene (double bonds) of the d-pulegone. This type of nucleophilic addition reaction occurs when the carbon-carbon bond has a carbonyl group adjacent to it [13]. The d-pulegone is a compound with double bonds adjacent to carbonyl group, also known as  $\alpha$ ,  $\beta$  unsaturated carbonyl compound. The  $\alpha$  and  $\beta$  signs indicate the distance of the double bonds from the carbonyl group, where the  $\alpha$  carbon is the carbon atom next to carbonyl group while the  $\beta$  carbon is the one further down the chain of the structure.

The nitrogen of the diethylamine protonates and became positively charged by interacting with hydrogen sulfite gas. The hydrogen sulfite ion produced, attacked the double bonds of d-pulegone by donating an electron to form single bond. An acidified hydrochloric acid is added, prevents the further protonation of diethylamine. The alkene bond being a planar (flat) exposed the carbon atom to a nucleophilic attacked either from above or below the plane of d-pulegone compound [14].

### **IV. CONCLUSION**

In this research work, a variety of analytical experiments and instruments were applied to purify, separate and identify the actual constituents using techniques such as TGA and ATR-FTIR analyses. The results provided by the experimental methodologies and instrumental analysis yielded results that satisfied the setting aim of the research work. In the TGA analysis, the result obtained showed the inflection point of pennyroyal oil (157 °C) falls within the range of inflection point of most the essential oils such Eugenol (162 °C), cinnamon (156 °C) etc. In the infrared analysis, slight shift of structural bonds was observed, which indicates chemical reactions and synthesis re-arrangement of reacting species is occurring. Further experiments and analytical processes are required to purify, separate and select the starting compounds and synthesize products such as column chromatography etc.

### **ACKNOWLEDGEMENT**

The authors gratefully acknowledge Dr Stephen Barton for supervising this work patiently.

## REFERENCES

- [1]. Asbahani, E.A. Miladi, A.K. and Elaissari, A. (2015) "Essential oil from extraction to encapsulation," J. Pharm., 483, pp. 220-243.
- [2]. Tongnuanchan, P. and Benjakul S., (2014) "Essential Oils: Extraction, Bioactivities, and Their Uses for Food," J. Food. Sci., 79, pp. R1231-R1249.
- [3]. Vian, A.M. Fernandez, X. Visinoni, F. and Chemat, F. (2008) "Microwave hydro-diffusion and Gravity, A New Technique for Extraction of Essential Oils," J. Chromatography, 1190, pp. 14-17
- [4]. Burt, S. (2004) "Essential Oil: Their antibacterial Properties and Potential Application in Food – A Review," J. Food. Microbiology, 94, pp. 223-253,
- [5]. Sandasi, M. Kamato, G.P.P. Branska, M. and Viljoen, A.M. (2010) "Application of Vibrational Spectroscopy in the Quality Assessment of Buchu Oil Obtained from Two Commercial Important *Agathosma* species (Rutaceae)," South Africa J. Botany, 76, pp. 692-700
- [6]. Chisowa, E. (1997) "Chemical Composition of Essential Oils of Three *Eucalyptus* Species in Zambia," J. Essential Oils Res., 9, pp. 653-655
- [7]. Gilles, M. Zhao, J. and Agboola, S. (2010) "Chemical Composition and Microbial Properties of Essential Oils of Three Australian *Eucalyptus* Species," Food Chemistry, 199, pp. 731-737
- [8]. Petrakis, A.E. Kimbaris, C.A. and Polissious, G.M. (2009) "Quantitative Determination of Pulegone in Pennyroyal oil by FT-IR Spectroscopy," J. Agriculture and Food Chem, 57, pp. 10044-10048
- [9]. Cremasco, A.M. and Nazarene, P.B. (2011) "Thermo gravimetric analysis of Long Pepper (*Piper hispidinervium* C. DC.) Essential Oil," Acta Amazonica, 15, pp. 5967
- [10]. Hazra, A. Alexander, K. Dollimore, D. and Riga, A. (2004) "Characterization of Some Essential Oils and Their Key Components: Thermo analytical techniques," J. Thermal Analysis and Calometry, 75, pp. 317-330
- [11]. Dewick, M.P. (2006) Essentials of Organic Chemistry, 1<sup>st</sup> ed., John Wiley & Sons: Chichester, pp.234.
- [12]. Caron, S. (2011) Practical Synthetic Organic Chemistry: Reactions, Principles and Techniques, 1<sup>st</sup> ed., Wiley: Hoboken, pp.75.
- [13]. Solomons, T.W. and Fryhle, B.C. (2011) Organic Chemistry, 10<sup>th</sup> ed., John Wiley & Sons: Asia, pp.187-890.
- [14]. Wade, L.G. (2010) Organic Chemistry: Pearson International Edition, 7<sup>th</sup> ed., Prentice Hall: London, pp. 379