

Evaluation of *Azotobacter* Isolates of Fodder Grasses and its Effect on Growth Parameters of Sweet Corn Crop

A. R. Potdar*, T.K. Narute, and A. R. Kedar

Department of Plant Pathology and Agricultural Microbiology

College of Agriculture, Pune

Mahatma Phule Krishi Vidyapeeth, Maharashtra State, India

*Corresponding author

ABSTRACT

A field experiment was conducted to evaluate the effect of *Azotobacter* isolates collected from rhizospheric soil of fodder grasses viz., Napier grass, Marvel grass, Anjan grass, Stylo grass, Johnson grass, Fodder jowar and Fodder maize as a seven treatments. The morphological, biochemical and cultural characteristics of individual isolate was studied. The application of lignite based biofertilizer of individual *Azotobacter* isolates as a seed treatment to the sweet corn seeds showed increased trend in various growth parameters in which the yield of sweet corn obtained due to treatment of *Azotobacter* isolated from rhizospheric soil of Napier grass found significantly higher (265.87 g/plant) than rest of treatments. Hence, the *Azotobacter* strain of Napier grass can be commercialized as an efficient biofertilizer strain of *Azotobacter*.

Keywords- *Azotobacter*, Fodder grasses, Sweet corn, Yield

Date of Submission: 08-07-2022

Date of acceptance: 22-07-2022

I. Introduction

Maize (*Zea mays* L.), is considered as Queen of Cereals. It is the world's third most important crop after wheat and rice. It has global importance in world agriculture. It contributes about 20 % of world's total cereal production. It is one of the most versatile crops in nature, which can be grown over wide range of climatic conditions. India produces about 2% of world's total maize production. The total maize production of India in 2017-18 was 21.93 MMT (Anonymous, 2017). There are six major types of maize as dent corn, flour corn, flint corn, pod corn, pop corn and sweet corn. Sweet corn (*Zea mays* var. *saccharata*) is also called as sugar corn and pole corn. It is a cereal with high sugar content. The sweet corn grain contains 16% carbohydrates, 6% protein, 3.22% sugar, 2.7% dietary fiber, 7% total fat, vitamin C 11%, vitamin B-6 5%, potassium 270mg/100g, iron 2%, magnesium 9%.

Sweet corn requires more amount of nutrients hence it is also called as heavy feeder. Nitrogen (N) is a major macronutrient which affects growth and yield of sweet corn crop. To fulfill the nutrient requirement of crop farmers are using higher doses of chemical fertilizers. The applications of heavy amount of chemical fertilizers are deteriorating soil and environment. Hence to overcome this problem use of biofertilizers is one of the ways in organic farming to increase crop yield without damaging the environment and soil health.

Azotobacter is the non-symbiotic, free living, aerobic, heterotrophic bacterium. It fixes atmospheric nitrogen in the form of ammonium ions in the soil, it also produces plant growth promoting hormones like gibberellins, cytokinin and auxin (Azcon *et al.*, 1975; Kukreja *et al.*, 2004), ammonia, vitamins and growth promoting substances which are responsible for seed germination (J Gonzalez-Lopez *et al.*, 1986), protection against root pathogens (Varma *et al.*, 2001; Sindhu *et al.*, 2009). Bacterium fixes an average of 10-15 kg N/ha per year (Henzell and Norris, 1962). It found in soil and rhizosphere of many crops. Its population varies from negligible to 10⁴/g of soil. It is founded that it fixes about 10µg of nitrogen by consuming 1 g of glucose. It is able to grow at a pH range of 4.8–8.5 and fixes nitrogen at optimum pH of 7.0–7.5 (Dilworth *et al.*, 1988)

Sweet corn is one of the hosts for *Azotobacter*. Many times the Nitrogen fixing bacteria are isolated from different plants including vegetables, cereals, pulses, fruit plants for their efficiency test. However the fodder grasses are not considered so far for isolating such microbes. The fodder grasses thrive well in all adverse climatic conditions. Their rhizospheric soil play key role in nutrient management. It contains all beneficial microbes which help grasses to survive in all climatic condition, Hence the present research was planned to isolate *Azotobacter* from rhizospheric soil of fodder grasses, their morphological study and nitrogen fixing efficiency of individual *Azotobacter* isolate.

II. Materials and methods

The present investigation was undertaken for the evaluation of *Azotobacter* isolates of fodder grasses and its effect on growth parameters, nutrient uptake and yield of sweet corn var. Madhu-5. The field trial was conducted at Department of Plant Pathology and Microbiology, College of Agriculture, Pune-411005 during 2018-19. The live seven fodder grass samples viz., Napier grass, Marvel grass, Anjan grass, Stylo grass, Johnson grass, Fodder jowar and Fodder maize were collected from All India Co-ordinated Research Project on Forage Crops and Utilization, central campus Mahatma Phule Krishi Vidyapeeth, Rahuri and Agronomy field, College of Agriculture, Pune. The eight treatments were replicated thrice and are detailed under table 1.

The fresh rhizospheric soil sample of individual grass was collected for isolation of *Azotobacter* in laboratory. The isolation of *Azotobacter* was done on Jensen's media by dilution and plating method. The isolated *Azotobacter* plates were kept in BOD incubator at $28 \pm 2^{\circ}\text{C}$ temperature for 3 days. After incubation period the typical colony growth was observed and their purity was verified by simple staining and Koperloff and Berman's modified method of Gram staining (Salle, 1967). Slants of Jensen's media were prepared and the pure culture was maintained by periodical sub culturing on Jensen's agar slants. The morphological, cultural and biochemical characters of individual *Azotobacter* isolates was studied by following standard procedures.

By using individual *Azotobacter* isolates, seven lignite based biofertilizer were prepared which then used for seed treatment to sweet corn seeds. The experimental plot was ploughed to a medium depth with clod crushing, ridges and furrows were opened at a distance of 60 cm. The initial soil samples were collected for analysis of available major nutrients; nitrogen (N), phosphorous (P) and potassium (K). The recommended FYM was added @ 15-20 tones/ha. and irrigated. After achieving the field capacity, the biofertilizer treated seeds were sown in three replications. Immediately, after sowing, field was irrigated. The recommended dose of fertilizer of sweet corn crop is 120:60:40 kg/ha. (N:P:K). Only basal dose of full potassium and phosphorus was given as per the recommended dose except control plot through the straight fertilizers i.e. Single Super Phosphate and Murate of Potash. Nitrogenous fertilizer was not given to the any treatment plot to assess the performance of nitrogen fixing efficiency of individual *Azotobacter* isolate alone. The subsequent irrigations were given as per the irrigation schedule. The moisture of field was maintained throughout the growing period of crop. Thinning was carried out after 10 days of sowing to maintain plant population and healthy seedlings. Hand weeding carried out to make weed free plot.

The observation on seed germination, height of plant, number of leaves, length of leaves, width of leaves, number of cobs, cob yield per plant, total dry matter weight of plant were recorded manually on five randomly selected representative plants from each plot of each replication separately as well as yield and yield attributing characters were recorded as per the standard method. Yield attributes were also recorded at physiological maturity stage. The data obtained from different observations were calculated as per Randomized Block Design (RBD) by using the standard statistical methods (Panse and Sukhatme, 1967) for its statistical significance.

III. Results and discussion

Morphological, cultural and biochemical characters of individual *Azotobacter* isolate: An individual *Azotobacter* isolate showed rod to spherical shape, non acid fast, non spore former, milky white to white colonies with striation and cyst formation with dark brown to black pigmentations and were identified as *Azotobacter* spp. It showed positive result to catalase test. The colonies were circular, smooth, undulating and convex.

Plant growth parameters and yield attributes: The data was recorded at 30, 60 and 90 days after sowing with respect to different growth parameters. The data in table 1 analyzed by using standard statistical method. Among the different treatments highest seed germination percentage was recorded in treatment T₁ with *Azotobacter* isolated from rhizospheric soil of Napier grass (89.47%) which is followed by treatment T₅ with *Azotobacter* isolated from rhizospheric soil of Johnson grass (78.07%). The maximum plant height (144.23 cm), number of leaves (12.60), length of leaf (95.89 cm), width of leaf (10.01) and number of cobs (2.00) were observed in treatment T₁. Maximum cob yield (265.87 g plant⁻¹) was recorded in treatment T₁ which is followed by treatment T₅ (200.40 g plant⁻¹). The Significant highest total dry matter weight was recorded in treatment T₁ (37.67 g plant⁻¹). The significant least growth was observed in treatment T₈ which was control.

IX. Conclusion

The explosions of Indian population enhance the demand of cereals, pulses, fruits etc. To fulfill this demand of increased population farmers are using excessive amount of fertilizers which directly affecting the soil health and environment, hence to overcome this, we have to go for organic farming with biofertilizers.

Mostly *Azotobacter* biofertilizers are prepared from different *Azotobacter* isolates isolated from vegetables, cereals, pulses, fruit plants. However the fodder grasses are not considered so far for isolating such microbes. The present research revealed significant results from which it can be concluded that

- The rhizospheric soil of Napier grass along with other six grasses has the efficient *Azotobacter* strains which can be used to produce biofertilizers of *Azotobacter*.
- Seed treatment of biofertilizer prepared from *Azotobacter* isolate of Napier grass to the sweet corn seeds resulted in increase in plant growth parameters and yield over control.
- The yield obtained from treatment T₁ with *Azotobacter* isolated from rhizospheric soil of Napier grass (265.87 g/plant) was followed by treatment T₅ of *Azotobacter* isolated from rhizospheric soil of Johnson grass (200.40 g/plant).
- The *Azotobacter* strain of Napier grass can be used to produce biofertilizers of *Azotobacter*.

References

- [1]. Anonymous., 2017c. National Collateral Management Services Limited (NCoMM) Special report: September 2017).
- [2]. Azcon, R. and Barea, J. M. 1975. Synthesis of auxins, gibberellins and cytokinins by *Azotobacter vinelandii* and *Azotobacter beijerinckii* related to effects produced on tomato plants. *Plant and Soil*. 43: 609–619.
- [3]. Dilworth, M.J., Eady, R.R. and Eldridge, M.E. 1998. The Vanadium nitrogenase of *Azotobacter chroococcum* Reduction of acetylene and ethylene to ethane. *Biochem. J.* 249: 745–751.
- [4]. Gonzalez-Lopez, J., Salmeron, V., Martine Toledo, M.V., Ballesteros, F. and Ramos Cormenzana, A. 1986. Production of auxins, gibberellins and cytokinins by *Azotobacter vinelandii* ATCC 12837 in chemically-defined media and dialized soil media, *Soil Biol. Biochem.* 18: 119–120.
- [5]. Henzell, E.P. and Norris, D. O. 1962. Processes by which nitrogen is added to the soil or plant system. In : Commonwealth Bureau Pasture and Field Crops, *Bull.* 46:1-25.
- [6]. Kukreja, K., Suneja, S., Goyal, S. and Narula, N. 2004. Phytohormone production by *Azotobacter*—a review, *Agric. Rev.* 25 (1): 70–75.
- [7]. Panse, V.G.S. and Sukhatme, V., 1967. Statistical Methods for Agricultural Workers. Indian Council of Agricultural Research. New Delhi. pp. 145-156.
- [8]. Salle, A.J. 1967. Laboratory manual of Fundamental Principles of bacteriology. McGraw Hill Book., New York.
- [9]. Sindhu, S.S., Rakshiya, Y.S. and Sahu, G. 2009. Biological control of soil-borne plant pathogens with rhizosphere bacteria. *Pest Technol.* 3: 10–21.
- [10]. Varma, A., Kukreja, K., Pathak, D.V., Suneja, S. and Nanda, N. 2001. *In vitro* production of plant growth regulators by *A. chroococcum*. *Indian Journal of Microbiology.* 41: 305-307.

Table 1. Effect of *Azotobacter* isolates of different fodder grasses on growth parameters and yield of sweet corn var. Madhu-5

Sr. No	Treatment	Treatment details	Germination (%)	Plant height (cm)	No. of leaves	Length of leaf (cm)	Width of leaf (cm)	No of cobs per plant	Cob yield per plant	Total dry matter
1	T ₁	<i>Azotobacter</i> isolated from rhizospheric soil of Napier grass (<i>Phule Gunvant</i>)	89.47	144.73	12.60	95.89	10.01	2.00	265.87	37.67
2	T ₂	<i>Azotobacter</i> isolated from rhizospheric soil of Marvel grass (<i>Phule Goverdhan</i>)	77.19	119.07	10.00	82.84	8.95	1.07	188.07	31.73
3	T ₃	<i>Azotobacter</i> isolated from rhizospheric soil of Anjan grass (<i>Phule Madras Anajan-1</i>)	78.07	140.80	10.67	80.27	8.86	1.00	163.87	27.37
4	T ₄	<i>Azotobacter</i> isolated from rhizospheric soil of Stylo grass (<i>Phule Kranti</i>)	73.68	140.60	10.07	78.27	8.75	1.00	169.20	30.97
5	T ₅	<i>Azotobacter</i> isolated from rhizospheric soil of Johnson grass (Local)	78.07	141.73	11.33	88.51	9.27	1.13	200.40	36.33
6	T ₆	<i>Azotobacter</i> isolated from rhizospheric soil of Fodder Maize (African tall)	72.81	124.27	10.47	83.60	9.06	1.07	157.47	31.87
7	T ₇	<i>Azotobacter</i> isolated from rhizospheric soil of Fodder Jowar (<i>Ruchira</i>)	64.91	139.20	10.37	84.31	9.02	1.00	183.20	28.50
8	T ₈	Absolute control (No culture or fertilizer)	53.51	117.33	9.20	71.64	7.94	1.00	121.67	25.00
SE±			5.06	5.74	0.51	2.85	0.27	0.09	13.95	1.52
CD(0.05)			15.51	17.59	1.57	8.74	0.84	0.28	42.73	4.67